

PEROXigard®

Wipes

One-Step^{††} Disinfectant Cleaner & Deodorizer

FOR USE IN LIFE SCIENCES

This product is a one-step^{††} disinfectant cleaner and deodorant designed for general cleaning, disinfecting, and deodorizing of hard, non-porous surfaces. It cleans quickly by removing urine, fecal matter, blood, food residue, dead skin, dirt, and other common soils found in lab animal research facilities, life science laboratories, small and large animal facilities, vivariums, animal housing, transportation vehicles, fish facilities, breeding establishments and operating rooms.

This product is designed for use on hard, non-porous surfaces such as: chrome, glass, glazed ceramic, glazed porcelain, glazed tile, plastic surfaces, stainless steel, and vinyl associated with high touch surfaces including countertops, biosafety cabinets, animal equipment, handheld devices, computer screens, cages, carts, and handling equipment.

Claims

BACTERICIDAL IN 1 MINUTE:

Acinetobacter baumannii (Multi-Drug Resistant (MDR))
Bordetella bronchiseptica
Bordetella pertussis
Burkholderia cepacia
Chlamydia psittaci[†]
Corynebacterium bovis[†]
Enterobacter aerogenes
Enterobacter cloacae
 (New Delhi metallo-beta-lactamase 1 (NDM 1))
Enterococcus faecalis
 (Vancomycin Resistant (VRE))
Enterococcus faecium
Escherichia coli
Escherichia coli O157:H7
Escherichia coli (extended spectrum beta-lactamase (ESBL))
Escherichia coli (New Delhi metallo-beta-lactamase 1(NDM-1))
Klebsiella pneumoniae
Klebsiella pneumoniae
 (Carbapenem Resistant (CRE))
Klebsiella pneumoniae (extended spectrum beta-lactamase (ESBL))
Klebsiella pneumoniae
 (New Delhi metallo-beta-lactamase 1 (NDM-1))
Legionella pneumoniae

Listeria monocytogenes
Pasteurella multocida
Pseudomonas aeruginosa
Pseudomonas aeruginosa
 (metallo-beta-lactamase (MBL) positive)
Salmonella enterica
Serratia marcescens
Shigella dysenteriae
Staphylococcus aureus
Staphylococcus aureus
 (Community Acquired Methicillin Resistant Genotype USA 400 (CA-MRSA 400))
Staphylococcus aureus
 (Methicillin Resistant (MRSA))
Staphylococcus aureus
 (Vancomycin Intermediate Resistant (VISA))
Staphylococcus aureus
 (Vancomycin Resistant (VRSA))
Staphylococcus epidermidis
 (Methicillin Resistant (MRSE))
Staphylococcus pseudintermedius
Stenotrophomonas (Pseudomonas) maltophilia
Streptococcus pneumoniae
 (Penicillin Resistant (PRSP))
Streptococcus pyogenes

Reference Sheet

KEEP OUT OF REACH OF CHILDREN

Manufactured by:

Virox Technologies Inc.
 2770 Coventry Road
 Oakville, ON Canada L6H 6R1
 1-800-387-7578 | virox.com



*VIRUCIDAL IN 1 MINUTE:

Adenovirus Type 8
 Avian Influenza A Virus (H3N2)
 Avian Influenza A Virus (H7N9)
 Avian Reovirus
 Bovine Viral Diarrhea Virus
 Canine Distemper Virus
 Canine Influenza A Virus
 Cytomegalovirus
 Duck Hepatitis B Virus
 Equine Herpes Virus Type 1
 Equine Influenza A Virus (H3N8) (EIV)
 Feline Calicivirus
 Feline Herpes Virus (Feline Rhinotracheitis)
 Feline Infectious Peritonitis Virus (FIPV)
 Feline Leukemia Virus
 Feline Panleukopenia Virus
 Hepatitis B Virus
 Hepatitis C Virus
 Herpes Simplex Virus Type 1
 Herpes Simplex Virus Type 2
 HIV Type 1 (Strain HTLV III_B)
 Human Coronavirus
 Influenza A Virus (Strain A/8/68)
 Influenza B Virus
 Marek's Disease Virus
 Minute Virus of Mice (Murine Parvovirus)
 Mouse Hepatitis Virus
 Murine Norovirus
 Newcastle Disease Virus
 Norovirus
 Pandemic Influenza A Virus (H1N1)
 Parainfluenza Virus Type 3
 Poliovirus Type 1 (Strain Chat)
 Porcine Epidemic Diarrhea Virus
 Porcine Respiratory & Reproductive Syndrome Virus
 Porcine Rotavirus
 Pseudorabies Virus
 Respiratory Syncytial Virus (RSV)

Rhinovirus Type-37
 Rotavirus (Strain WA)
 SARS-Related Coronavirus 2 (Human Coronavirus SARS-CoV-2 (the virus that causes COVID-19*))
 Vaccinia Virus
 Vesicular Stomatitis Virus

 Meets surface disinfection recommendations under the OSHA Bloodborne Pathogen Standard for *HBV, *HCV & *HIV-1

TUBERCULOCIDAL IN 1 MINUTE:

Mycobacterium bovis (BCG)

FUNGICIDAL IN 1 MINUTE:

Candida albicans
Microsporium canis
Trichophyton interdigitale
 (formerly known as *Trichophyton mentagrophytes*)

NON-FOOD CONTACT SANITIZER IN 10 SECONDS:

Acinetobacter baumannii
Campylobacter jejuni
Enterobacter aerogenes
Enterococcus faecalis
 (Vancomycin Resistant (VRE))
Escherichia coli O157:H7
Listeria monocytogenes
Pseudomonas aeruginosa
Salmonella enterica
Staphylococcus aureus
Staphylococcus aureus
 (Methicillin Resistant (MRSA))

MOLD/MILDEW (MILDEWCIDAL ACTIVITY) IN 4 MINUTES:

Aspergillus brasiliensis
 (formerly known as *Aspergillus niger*)

Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**Not for personal cleansing. This is not a baby wipe.
Do not flush in toilet.**

For Use as a One-Step* Cleaner/Disinfectant:

1. Pre-clean visibly soiled areas.
2. Pull towelette from canister and wipe hard, non-porous surfaces.
3. Allow surfaces to remain visibly wet for 1 minute.[¶]
4. Allow to air dry.

For Use to Clean and Disinfect Life Science Laboratory Surfaces, Instruments, and Equipment:

1. Pre-clean visibly soiled areas.
2. Pull towelette from canister and wipe hard, non-porous surfaces.
3. Allow surfaces to remain visibly wet for 1 minute.[¶]
4. If streaking is observed, wipe with a clean, damp cloth or microfiber cloth.

For Use as a Cleaner/Disinfectant in Animal Housing Facilities:

For cleaning and disinfecting the following hard, non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and fecal matter from floors, walls and surfaces of facilities occupied by animals.
3. Empty all feeding or watering appliances.
4. Thoroughly clean all surfaces with soap and detergent and rinse with potable water. Pull towelette from canister and wipe hard, non-porous surfaces.
5. All surfaces must remain visibly wet for 1 minute.[¶]
6. Allow to air dry.
7. Thoroughly scrub all treated automatic feeders and waterers with soap or detergent, and rinse with potable water before reuse.

For Cleaning and Disinfection of Vehicle Cabins:

To disinfect hard, non-porous surfaces of vehicle cabins:

1. Pre-clean visibly soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. Allow surface to remain visibly wet for 1 minute.[¶]
4. Allow to air dry.

To Kill Mold and Mildew:

1. Pre-clean surfaces.
2. Pull towelette from canister and wipe product onto hard, non-porous surfaces until completely wet.
3. Allow the surface to remain visibly wet for 4 minutes.
4. Allow to air dry.

To Disinfect Non-Critical[¶], Pre-cleaned Instruments:

1. Instruments must be thoroughly pre-cleaned to remove excess organic debris, rinsed, and rough dried.
2. Wipe all hard, non-porous surfaces of instruments with this product until thoroughly wet.
3. Allow surfaces to remain visibly wet for 1 minute.[¶]
4. Allow to air dry.

For Use as a Non-Food Contact Sanitizer:

1. Pre-clean visibly soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. Allow the surface to remain visibly wet for 10 seconds.
4. Allow to air dry.

For Use as a Cleaner and/or Deodorizer:

1. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
2. Allow to air dry.

To Kill *Corynebacterium bovis* (*C.bovis*):

1. Pre-clean surfaces by following the cleaner instructions listed above.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. Allow surfaces to remain visibly wet for 2 minutes.
4. Allow to air dry.

To Kill *Chlamydia psittaci*:

1. Pre-clean surfaces by following the cleaner instructions listed above.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. Allow surfaces to remain visibly wet for 1 minute.
4. Allow to air dry.

***KILLS HIV-1, HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings and other settings in which there is an expected likelihood of soiling of hard, non-porous surfaces / objects with blood / body fluids, and in which the surfaces / objects likely to be soiled with blood / body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1), Hepatitis B Virus (HBV), or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV OR HCV ON SURFACES / OBJECTS SOILED WITH BLOOD / BODY FLUIDS:

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood / body fluids must be thoroughly cleaned from surfaces / objects before application of this product.

Contact Time: Allow surface to remain visibly wet for 1 minute[¶] to kill HIV-1, HBV and HCV.

Disposal of Infectious Material: Cleaning materials used that may contain blood / body fluids should be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

USER SHOULD READ S.D.S. BEFORE USING PRODUCT
SDS is available online at www.virox.com

DISCLAIMER: You must read and follow the instructions on the label for our product and use the product for the particular applications specified on the product label. To the full extent permitted under applicable law, Virox Technologies Inc. shall not be liable for any damages whatsoever, arising out of or otherwise in connection with any failure to follow those instructions or any use of its products which is not specified on the product's label. The instructions on the product label are not a warranty, express or implied, and Virox Technologies, Inc. expressly disclaims any implied warranties, including any warranty of fitness for a particular purpose. Under applicable law, some of the above waivers or disclaimers of warranty, or exclusions of liability for certain types of damages may not apply to you. In such cases, our liability will be limited to the fullest extent permitted by applicable law.

*Hard, non-porous surfaces

† For use against *Corynebacterium bovis* and *Chlamydia psittaci* please see the specific use directions.

¶ Non-Critical medical devices are items that come in contact only with intact skin.

¶¶ Use a 2 minute contact time against *Corynebacterium bovis*.

†† When used according to disinfection use directions.

™ Accelerated Hydrogen Peroxide® and Design are trademarks of Diversey, Inc. | "Accelerated" does not imply increased product efficacy.